

Frugivorous Psittacine Crop Milk

DESCRIPTION

It is formulated to hand-rear frugivorous psittacine birds during the first phase of the early-age period from hatching until the 7th-8th day of life (depending on the species). It is a highly concentrated and digestible formula. It is formulated to constitute 100 % of the chick's diet during that period.

This product is especially recommended for breeders who incubate eggs artificially. It provides vigorous growth similar to those achieved by the best breeding pairs. This vigour promotes health and minimizes the risk of contracting diseases.

FEEDING RECOMMENDATIONS

On day one, the concentration should be increased from approximately 1 % in the first feeding bout to 20 % in the last one.

From day 3, use a concentration of 25 %.

Psittacus Frugivorous Psittacine Crop Milk has to be switched progressively to **Psittacus Frugivorous Neonatal Hand Feeding**.

Leftover formula must be discarded after each feeding bout.

After opening, store in an air-tight container in a cool dry place.

While refrigeration is not advised, freezing is possible if it is kept hermetically sealed.

PREPARATION

It should be prepared with warm water between 37-39 °C.

DAILY INTAKE

It is important to offer the right amount to the chicks divided in 7-8 feeding bouts during the first few days.

NUTRITIONAL INFORMATION

COMPOSITION:

Oil seeds, oil fruits and products derived thereof, fructose, vegetable oils (high oleic sunflower, crude palm, refined linseed), dried whole egg, minerals and products derived thereof, inulin (*Cichorium intybus*), marigold meal (*Tagetes erecta*).

ADDITIVES:

Vitamins: 33a672a Vitamin A: 4,000 IU/kg; 3a671 Vitamin D3: 1,300 IU/kg; 3a700 Vitamin E: 330 mg/kg; 3a300 Vitamin C: 200 mg/kg; 3a710 Vitamin K3: 5 mg/kg; 3a820 Vitamin B1: 25 mg/kg; 3a825i Vitamin B2: 20 mg/kg; 3a831 Vitamin B6/Pyridoxine hydrochloride: 25 mg/kg; Vitamin B12/Cyanocobalamin: 100 µg/kg; 3a841 Calcium D-pantothenate: 20 mg/kg; 3a314 Niacin: 125 mg/kg; 3a316 Folic acid: 5 mg/kg; 3a880 Biotin: 800 µg/kg; 3a890 Choline chloride: 1,600 mg/kg; 3a160 (a) Beta-carotene: 13 mg/kg; 3a920 Betaine anhydrous: 250 mg/kg. **Trace elements:** 3b106 Iron (iron (II) chelate of amino acids hydrate): 20 mg/kg; 3b406 Copper (copper (II) chelate of amino acids hydrate): 1 mg/kg; 3b504 Manganese (manganese chelate of amino acids hydrate): 50 mg/kg; 3b606 Zinc (zinc chelate of amino acids hydrate): 20 mg/kg; 3b202 Iodine (calcium iodate anhydrous): 0.8 mg/kg; 3b814 Selenium (hydroxy-analogue of selenomethionine): 210 µg/kg. **Antioxidants:** 1b306 (ii) Extract of vegetable oils rich in tocopherols (rich in delta-tocopherol): 226 mg/kg; 1b304 Ascorbyl palmitate: 50 mg/kg. **Flavourings.**

ANALYTICAL CONSTITUENTS:

Moisture 6.0 %, Crude protein 39.0 %, Crude fat 29.5 %, Crude fibre 2.8 %, Crude ash 6.7 %.

PACKAGING

500 g bag.



500 g

